

Appendix B

Mostly linear algebra: a brief review

We assume that you have taken an introductory linear algebra course in addition to multivariable calculus. The following is only a quick discussion of some terminology and notation used in this book. We do not give proofs of all results.

B.1 Polynomials and sequences

A *polynomial* is an expression of the form

$$p(z) = a_0 + a_1z + \cdots + a_nz^n. \quad (\text{B.1.1})$$

In this book, the coefficients a_0, \dots, a_n will always be complex numbers, and z will usually be complex, so that p is a function from \mathbb{C} to \mathbb{C} . We will then call p either a polynomial or a complex polynomial. If all the coefficients are real numbers, we will call p a *real polynomial*; if they are rational, we will call it a *rational polynomial*. (There are other possibilities: in coding theory, for instance, the coefficients of a polynomial may be elements of a finite field.) A real polynomial can take on nonreal values. For example, let $p(z) := 2z^2$ for all $z \in \mathbb{C}$. Then p is a real polynomial but $p(1+i)$ is not a real number.

A real polynomial on a nonempty subset $S \subset \mathbb{C}$ is the restriction of a real polynomial to S . A complex polynomial on S is the restriction of a complex polynomial to S .

We denote by $\mathbb{C}[t]$ the collection of all complex polynomials and by $\mathbb{R}[t]$ the collection of all real polynomials.

Since the identity function $i: \mathbb{C} \rightarrow \mathbb{C}$ defined by $i(z) := z$ is continuous, Proposition 1.5.14 implies that every polynomial is continuous on all of \mathbb{C} . Hence, for any nonempty subset S of \mathbb{C} , Proposition 1.5.12 implies that every polynomial function on S is continuous on all of S .

Recall that in calculus, a sequence in \mathbb{R} is a function $f: \mathbb{N} \rightarrow \mathbb{R}$.

Definition B.1.1 (Sequence). Let X be any nonempty set. A *sequence in X* is a function

$$f : \mathbb{N} \rightarrow X.$$

If f is a sequence in X and $x_n := f(n)$ for each $n \in \mathbb{N}$, we will often denote f by (x_n) . This notation is less cumbersome and more intuitive than the functional notation f . So, for instance, “let (x_n) be a sequence in X ” means that (x_n) denotes a sequence

$$f : \mathbb{N} \rightarrow X, \quad \text{where } f(n) = x_n \quad \text{for all } n \in \mathbb{N}. \quad (\text{B.1.2})$$

We sometimes denote the sequence (x_n) by an infinite tuple (x_1, x_2, \dots) . Each x_i is a *term* of the sequence (x_n) . The k^{th} term of a sequence (x_n) is x_k .

If $k \neq l$, then x_k and x_l are said to be *different terms* of the sequence even if $x_k = x_l$. Hence, every sequence (x_n) always has infinitely many different terms, whereas the set $\{x_1, x_2, \dots\}$ that contains all the terms of the sequence may be finite. If M is a subset of a set X and (x_n) is a sequence in X , we write $(x_n) \subset M$ to mean that $x_n \in M$ for all n .

B.2 Vector spaces

Let \mathcal{K} be either \mathbb{R} or \mathbb{C} . A *vector space over \mathcal{K}* is a triple $(V, +, \bullet)$, where V is a nonempty set and $+: V \times V \rightarrow V$ and $\bullet: \mathcal{K} \times V \rightarrow V$ are functions satisfying certain properties. The elements of V are then called *vectors*, \mathcal{K} is called the *scalar field of V* , elements of \mathcal{K} are called *scalars*, the function $+$ is called the *addition operation*, and \bullet is called the *scalar multiplication operation*. For v, w in $V \times V$ and $\alpha \in \mathcal{K}$, we will write $v + w$ for $+(v, w)$ and we write αv for $\bullet(\alpha, v)$. We call $u + v$ the *sum of u and v* ; we call αv the *scalar multiplication of v by α* . The required properties of $+$ and \bullet are as follows:

1. *Additive identity:* There exists $\mathbf{0}_V \in V$ such that $\mathbf{0}_V + v = v$ for any $v \in V$.
2. *Additive inverse:* For any $v \in V$, there is $-v \in V$ such that $v + (-v) = \mathbf{0}$.
3. *Commutative law for addition:* $v + w = w + v$ for all $v, w \in V$.
4. *Associative law for addition:* For all $v_1, v_2, v_3 \in V$,

$$v_1 + (v_2 + v_3) = (v_1 + v_2) + v_3.$$

5. *Multiplicative identity:* For all $v \in V$, we have $1v = v$.
6. *Associative law for multiplication:* For all $\alpha, \beta \in \mathcal{K}$ and all $v \in V$, $\alpha(\beta v) = (\alpha\beta)v$.
7. *Distributive law for scalar addition:* For all scalars $\alpha, \beta \in \mathcal{K}$ and all $v \in V$, $(\alpha + \beta)v = \alpha v + \beta v$.
8. *Distributive law for vector addition:* For all $\alpha \in \mathcal{K}$ and $v, w \in V$, we have $\alpha(v + w) = \alpha v + \alpha w$.

Although the operations of addition and scalar multiplication defined on different vector spaces can be different, we will always use $+$ to denote the addition operation, and αv to denote the scalar multiplication of v by α .

A real vector space is one whose scalar field is \mathbb{R} ; a complex vector space is one whose scalar field is \mathbb{C} .

When it's clear or irrelevant what the addition and scalar multiplication operations are, we write simply " V is a vector space over \mathcal{K} ". When \mathcal{K} is clear or irrelevant, we may write " V is a vector space".

It is easy to show that every vector space V has a unique additive identity (the vector $\mathbf{0}_V$ in property 1 above). It is also called the zero vector. The zero vector of every vector space will be denoted by $\mathbf{0}$.

Let v_1, \dots, v_n be vectors in a vector space V . It follows from properties 3 and 4 above that the meaning of $v_1 + \dots + v_n$ does not depend on the order in which the v_k are written.

Definition B.2.1 (Vector subspace). Let V be a vector space over \mathcal{K} and let M be a nonempty subset of V . Then M is a *vector subspace* of V if for all m_1, m_2 in M and all $\alpha \in \mathcal{K}$, we have $m_1 + m_2 \in M$ and $\alpha m_1 \in M$ – i.e., M is closed under the operations of addition and scalar multiplication defined on V .

A vector subspace M is *nonzero* (or *nontrivial*) if $M \neq \{\mathbf{0}\}$. It is *trivial* if $M = \{\mathbf{0}\}$. Do not confuse the trivial vector subspace with the empty set; the empty set is not a vector subspace. A vector subspace M is *proper* if $M \neq V$.

Notation B.2.2 (Notation for \mathbb{R}^n and \mathbb{C}^n). Throughout this book, we will denote elements of \mathcal{K}^n (where \mathcal{K} is either \mathbb{R} or \mathbb{C}) either by a bold letter or by a column matrix whose entries are indexed non-bold versions of the same

letter: $\mathbf{x} := \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}$, $\mathbf{v} := \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}$, and so on (sometimes we also write such an element as the transpose of a line matrix). If \mathbf{x} is in \mathbb{R}^n , then x_1, \dots, x_n are real numbers; if \mathbf{x} is in \mathbb{C}^n , then x_1, \dots, x_n are complex numbers.

We write vectors in \mathbb{R}^n and \mathbb{C}^n as column matrices because (see Theorem B.5.2) every linear transformation from \mathcal{K}^n to \mathcal{K}^m is given by multiplication by a $m \times n$ matrix A ; writing these vectors in column form enables us to use the familiar $f(x) = y$ format, as we do in (B.5.4), for example. It would not make sense to write that equation as $\begin{bmatrix} 1 & 4 & -2 \\ 3 & 0 & 2 \end{bmatrix} (1, -1, -2)$.

Let V be a vector space over \mathcal{K} . Let A, B be nonempty subsets of V and let $v \in V, \alpha \in \mathcal{K}$. Then

$$v + A := \{v + a \mid a \in A\} \tag{B.2.1}$$

$$A + B := \{a + b \mid a \in A, b \in B\} \tag{B.2.2}$$

$$\alpha B := \{\alpha b \mid b \in B\}. \tag{B.2.3}$$

We also define $A + v$ to be $v + A$ and $-B := (-1)B$.

Hence, αB can be obtained from B by “stretching” or “shrinking” each vector in B by a factor of $|\alpha|$. Consequently, if B is an open ball in \mathbb{R}^2 with radius $r > 0$, then αB is an open ball in \mathbb{R}^2 with radius $|\alpha|r$. Although the above operations look like addition and multiplication of numbers, if A is a vector subspace of V , then $A = -A$ and A and $-A$ don’t cancel.

Definition B.2.3. (Balanced subset, convex subset). Let V be a vector space with scalar field \mathcal{K} .

1. $M \subset V$ is *balanced* if $tM \subset M$ whenever $t \in \mathcal{K}$ and $|t| \leq 1$.
2. $M \subset V$ is *convex* if whenever x, y are in M and $t \in [0, 1]$, then $tx + (1 - t)y$ is also in M .

Thus every vector subspace of V is convex and balanced. If M is a subset of \mathbb{R}^n , then we can restate the definition of convexity to say that M is convex if for any two points $\mathbf{x}, \mathbf{y} \in M$, the straight line segment joining \mathbf{x} to \mathbf{y} lies entirely in M .

Example B.2.4. Any convex, symmetric subset U containing $\mathbf{0}$ must be balanced. To see this, let t be a scalar with $0 < |t| \leq 1$. Then since U is convex and $\mathbf{0} \in U$, we have $|t|U = |t|U + (1 - |t|)\{\mathbf{0}\} \subset |t|U + (1 - |t|)U \subset U$. Hence, since U is symmetric, $\frac{t}{|t|}U \subset U$ so that

$$tU = |t| \left(\frac{t}{|t|} U \right) \subset |t|U \subset U.$$

If $t = 0$, then we also have $tU \subset U$, because $\mathbf{0} \in U$. Thus, U is balanced.

Product of vector spaces

Assume that X_1, \dots, X_n are vector spaces over a common scalar field \mathcal{K} . Let

$$\mathbf{X} := \{ (x_1, \dots, x_n) \mid x_j \in X_j, 1 \leq j \leq n \} = X_1 \times \dots \times X_n. \quad (\text{B.2.4})$$

For $\mathbf{x} := (x_1, \dots, x_n)$, $\mathbf{y} := (y_1, \dots, y_n)$ in \mathbf{X} and $\alpha \in \mathcal{K}$, define

$$\begin{aligned} \mathbf{x} + \mathbf{y} &:= (x_1 + y_1, \dots, x_n + y_n) \quad \text{and} \\ \alpha \mathbf{x} &:= (\alpha x_1, \dots, \alpha x_n). \end{aligned} \quad (\text{B.2.5})$$

If as usual we denote by $\mathbf{0}$ the zero vector in each X_j , it can be verified easily that \mathbf{X} is a vector space over \mathcal{K} with $(\mathbf{0}, \dots, \mathbf{0})$ as the zero vector. The vector space \mathbf{X} is called the *product* of the vector spaces X_1, \dots, X_n .

B.3 Linear independence and span

The definitions of *linear combination* and *linear independence* in the context of finite-dimensional vector spaces remain valid in infinite dimensions.