

2.10.15 a. Show that the mapping

$$F \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} e^x + e^y \\ e^x + e^{-y} \end{pmatrix} \text{ is locally invertible at every point } \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2.$$

b. If $F(\mathbf{a}) = \mathbf{b}$, what is the derivative of F^{-1} at \mathbf{b} ?

2.10.16 The matrix $A_0 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$ satisfies $A_0^3 = I$.

True or false? There exists a neighborhood $U \subset \text{Mat}(3, 3)$ of I and a continuously differentiable function $g : U \rightarrow \text{Mat}(3, 3)$ with $g(I) = A_0$ and $(g(A))^3 = A$ for all $A \in U$ (i.e., $g(A)$ is a cube root of A).

2.10.17 Prove theorem 2.10.2 (the inverse function theorem in one dimension).

2.11 REVIEW EXERCISES FOR CHAPTER 2

$$x + y - z = a$$

$$x + 2z = b$$

$$x + ay + z = b$$

Equations for exercise 2.1

$$\begin{bmatrix} 1 & 2 & 0 & 1 \\ 1 & 1 & 3 & 3 \\ 0 & 1 & 0 & 1 \\ 2 & 1 & 1 & 3 \end{bmatrix}$$

Matrix A of exercise 2.2

$$\begin{bmatrix} 1 & -1 & 3 & 0 & -2 \\ -2 & 2 & -6 & 0 & 4 \\ 0 & 2 & 5 & -1 & 0 \\ 2 & -6 & -4 & 2 & -4 \end{bmatrix}$$

Matrix A for exercise 2.4.

Exercise 2.4, part b: For example, for $k = 2$ we are asking about the system of equations

$$\begin{bmatrix} 1 & -1 \\ -2 & 2 \\ 0 & 2 \\ 2 & -6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ -6 \\ 5 \\ -4 \end{bmatrix}.$$

2.1 a. For what values of a and b does the system of linear equations shown in the margin have one solution? No solutions? Infinitely many solutions?

b. For what values of a and b is the matrix of coefficients invertible?

2.2 When A is the matrix at left, multiplication by what elementary matrix corresponds to

a. Exchanging the first and second rows of A ?

b. Multiplying the fourth row of A by 3?

c. Adding 2 times the third row of A to the first row of A ?

2.3 a. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Are the following statements true or false?

1. If $\ker T = \{\vec{0}\}$, and $T(\vec{y}) = \vec{b}$, then \vec{y} is the only solution to $T(\vec{x}) = \vec{b}$.

2. If \vec{y} is the only solution to $T(\vec{x}) = \vec{c}$, then for any $\vec{b} \in \mathbb{R}^m$, a solution exists to $T(\vec{x}) = \vec{b}$.

3. If $\vec{y} \in \mathbb{R}^n$ is a solution to $T(\vec{x}) = \vec{b}$, it is the only solution.

4. If for any $\vec{b} \in \mathbb{R}^m$ the equation $T(\vec{x}) = \vec{b}$ has a solution, then it is the only solution.

b. For any statements that are false, can one impose conditions on m and n that make them true?

2.4 a. Row reduce the matrix A in the margin.

b. Let \vec{v}_m , $m = 1, \dots, 5$ be the columns of A . What can you say about the systems of equations

$$[\vec{v}_1, \dots, \vec{v}_k] \begin{bmatrix} x_1 \\ \vdots \\ x_k \end{bmatrix} = \vec{v}_{k+1} \quad \text{for } k = 1, 2, 3, 4?$$

2.5 a. Let A be an invertible $n \times n$ matrix, B an invertible $m \times m$ matrix, C any $n \times m$ matrix, and $[0]$ the $m \times n$ zero matrix. Show that the $(n+m) \times (n+m)$ matrix $\begin{bmatrix} A & C \\ [0] & B \end{bmatrix}$ is invertible.

b. Find a formula for the inverse.

2.6 Exercise 2.2.11 asked you to show that using row reduction to solve n equations in n unknowns takes $n^3 + n^2/2 - n/2$ operations, where a single addition, multiplication, or division counts as one operation. How many operations are needed to compute the inverse of an $n \times n$ matrix A ? To perform the matrix multiplication $A^{-1}\vec{b}$?

2.7 a. For what values of a is the matrix $\begin{bmatrix} 1 & -1 & -1 \\ 0 & a & 1 \\ 2 & a+2 & a+2 \end{bmatrix}$ invertible?

b. For those values, compute the inverse.

2.8 Show that the following two statements are equivalent to saying that a set of vectors $\vec{v}_1, \dots, \vec{v}_k$ is linearly independent:

a. The only way to write the zero vector $\vec{0}$ as a linear combination of the \vec{v}_i is to use only zero coefficients.

b. None of the \vec{v}_i is a linear combination of the others.

2.9 a. Show that $\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ form an orthonormal basis of \mathbb{R}^2 .

b. Show that $\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} \sin \theta \\ -\cos \theta \end{bmatrix}$ form an orthonormal basis of \mathbb{R}^2 .

c. Show that any orthogonal 2×2 matrix gives either a reflection or a rotation: a reflection if its determinant is negative, a rotation if its determinant is positive.

2.10 a. For vectors in \mathbb{R}^2 , prove that the length squared of a vector is the sum of the squares of its coordinates, with respect to any orthonormal basis.

b. Prove the same thing for vectors in \mathbb{R}^3 .

c. Repeat for \mathbb{R}^n : show that if $\vec{v}_1, \dots, \vec{v}_n$ and $\vec{w}_1, \dots, \vec{w}_n$ are two orthonormal bases, and $a_1\vec{v}_1 + \dots + a_n\vec{v}_n = b_1\vec{w}_1 + \dots + b_n\vec{w}_n$, then

$$a_1^2 + \dots + a_n^2 = b_1^2 + \dots + b_n^2.$$

2.11 a. Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$. Are the elements I, A, A^2, A^3 linearly independent in $\text{Mat}(2, 2)$? What is the dimension of the subspace $V \subset \text{Mat}(2, 2)$ that they span? (Recall that $\text{Mat}(n, m)$ denotes the set of $n \times m$ matrices.)

b. Show that the set W of matrices $B \in \text{Mat}(2, 2)$ that satisfy $AB = BA$ is a subspace of $\text{Mat}(2, 2)$. What is its dimension?

c. Show that $V \subset W$. Are they equal?

2.12 Let $\vec{v}_1, \dots, \vec{v}_k$ be vectors in \mathbb{R}^n , and set $V = [\vec{v}_1, \dots, \vec{v}_k]$.

a. Show that the set $\vec{v}_1, \dots, \vec{v}_k$ is orthogonal if and only if $V^\top V$ is diagonal.

b. Show that the set $\vec{v}_1, \dots, \vec{v}_k$ is orthonormal if and only if $V^\top V = I_k$.

2.13 Find a basis for the image and the kernel of the matrices

$$A = \begin{bmatrix} 1 & 1 & 3 & 6 & 2 \\ 2 & -1 & 0 & 4 & 1 \\ 4 & 1 & 6 & 16 & 5 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 1 & 3 & 6 & 2 \\ 2 & -1 & 0 & 4 & 1 \end{bmatrix},$$

Exercise 2.14: For example, the polynomial

$$p = 2x - y + 3xy + 5y^2$$

corresponds to the point $\begin{pmatrix} 0 \\ 2 \\ -1 \\ 0 \\ 3 \\ 5 \end{pmatrix}$, so

$$xD_1p = x(2 + 3y) = 2x + 3xy$$

corresponds to $\begin{pmatrix} 0 \\ 2 \\ 0 \\ 0 \\ 3 \\ 0 \end{pmatrix}$.

Hint for exercise 2.15: You should use the fact that a polynomial p of degree d such that $p(n) = p'(n) = 0$ can be written $p(x) = (x - n)^2 q(x)$ for some polynomial q of degree $d - 2$.

Hint for exercise 2.16, part b: $\vec{v} = P\vec{v} + (\vec{v} - P\vec{v})$.

Exercise 2.17: Recall that \mathcal{C}^2 is the space of C^2 (twice continuously differentiable) functions.

and verify that the dimension formula is true.

2.14 Let P be the space of polynomials of degree at most 2 in the two variables x, y , identified to \mathbb{R}^6 by identifying $a_1 + a_2x + a_3y + a_4x^2 + a_5xy + a_6y^2$ with $\begin{pmatrix} a_1 \\ \vdots \\ a_6 \end{pmatrix}$.

- a. What are the matrices of the linear transformations $S, T : P \rightarrow P$

$$S(p) \begin{pmatrix} x \\ y \end{pmatrix} = xD_1p \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{and} \quad T(p) \begin{pmatrix} x \\ y \end{pmatrix} = yD_2p \begin{pmatrix} x \\ y \end{pmatrix}?$$

- b. What are the kernel and the image of the linear transformation

$$p \mapsto 2p - S(p) - T(p)?$$

2.15 Let $a_1, \dots, a_k, b_1, \dots, b_k$ be any $2k$ numbers. Show that there exists a unique polynomial p of degree at most $2k - 1$ such that $p(n) = a_n, p'(n) = b_n$ for all integers n with $1 \leq n \leq k$. In other words, show that the values of p and p' at $1, \dots, k$ determine p .

2.16 A square $n \times n$ matrix P such that $P^2 = P$ is called a *projection*.

a. Show that P is a projection if and only if $I - P$ is a projection. Show that if P is invertible, then P is the identity.

b. Let $V_1 = \text{img } P$ and $V_2 = \ker P$. Show that any vector $\vec{v} \in \mathbb{R}^n$ can be written uniquely $\vec{v} = \vec{v}_1 + \vec{v}_2$ with $\vec{v}_1 \in V_1$ and $\vec{v}_2 \in V_2$.

c. Show that there exist a basis $\vec{v}_1, \dots, \vec{v}_n$ of \mathbb{R}^n and a number $k \leq n$ such that

$$P\vec{v}_1 = \vec{v}_1, P\vec{v}_2 = \vec{v}_2, \dots, P\vec{v}_k = \vec{v}_k \quad \text{and}$$

$$P\vec{v}_{k+1} = \vec{0}, P\vec{v}_{k+2} = \vec{0}, \dots, P\vec{v}_n = \vec{0}.$$

*d. Show that if P_1 and P_2 are projections such that $P_1P_2 = [0]$, then $Q = P_1 + P_2 - (P_2P_1)$ is a projection, $\ker Q = \ker P_1 \cap \ker P_2$, and the image of Q is the space spanned by the image of P_1 and the image of P_2 .

2.17 Show that the transformation $T : \mathcal{C}^2(\mathbb{R}) \rightarrow \mathcal{C}(\mathbb{R})$ given by formula 2.6.8 in example 2.6.9 is a linear transformation.

2.18 Denote by $\mathcal{L}(\text{Mat}(n, n), \text{Mat}(n, n))$ the space of linear transformations from $\text{Mat}(n, n)$ to $\text{Mat}(n, n)$.

a. Show that $\mathcal{L}(\text{Mat}(n, n), \text{Mat}(n, n))$ is a vector space and that it is finite dimensional. What is its dimension?

b. Prove that for any $A \in \text{Mat}(n, n)$, the transformations

$$L_A, R_A : \text{Mat}(n, n) \rightarrow \text{Mat}(n, n)$$

given by $L_A(B) = AB, R_A(B) = BA$ are linear transformations.

c. Let $\mathcal{M}_L \subset \mathcal{L}(\text{Mat}(n, n), \text{Mat}(n, n))$ be the set of functions of the form L_A . Show that it is a subspace of $\mathcal{L}(\text{Mat}(n, n), \text{Mat}(n, n))$. What is its dimension?

d. Show that there are linear transformations $T : \text{Mat}(2, 2) \rightarrow \text{Mat}(2, 2)$ that cannot be written as $L_A + R_B$. Can you find an explicit one?

e. What are $|L_A|$ and $|R_A|$ in terms of $|A|$ and $|B|$?

2.19 Show that in a vector space of dimension n , more than n vectors are never linearly independent, and fewer than n vectors never span.

2.20 Suppose we use the same operator $T : P_2 \rightarrow P_2$ as in exercise 2.6.8, but choose instead to work with the basis

$$q_1(x) = x^2, \quad q_2(x) = x^2 + x, \quad q_3(x) = x^2 + x + 1.$$

Now what is the matrix $\Phi_{\{q\}}^{-1} \circ T \circ \Phi_{\{q\}}$?

2.21 Let $V, W \subset \mathbb{R}^n$ be two subspaces.

a. Show that $V \cap W$ is a subspace of \mathbb{R}^n .

b. Show that if $V \cup W$ is a subspace of \mathbb{R}^n , then either $V \subset W$ or $W \subset V$.

2.22 a. Find a global Lipschitz ratio for the derivative of the map F defined in the margin.

b. Do one step of Newton's method to solve $F\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) - \begin{pmatrix} .5 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, starting at $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$.

c. Can you be sure that Newton's method converges?

2.23 Using Newton's method, solve the equation $A^3 = \begin{bmatrix} 9 & 0 & 1 \\ 0 & 7 & 0 \\ 0 & 2 & 8 \end{bmatrix}$.

2.24 Consider the map $F : \text{Mat}(2, 2) \rightarrow \text{Mat}(2, 2)$ given by $F(A) = A^2 + A^{-1}$.

Set $A_0 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ and $B_0 = F(A_0)$, and define

$$U_r = \{B \in \text{Mat}(2, 2) \mid |B - B_0| < r\}.$$

Do there exist $r > 0$ and a differentiable mapping $G : U_r \rightarrow \text{Mat}(2, 2)$ such that $F(G(B)) = B$ for every $B \in U_r$?

2.25 a. Find a global Lipschitz ratio for the derivative of the mapping $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given in the margin.

b. Do one step of Newton's method to solve $F\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ starting at $\begin{pmatrix} 2 \\ 3 \end{pmatrix}$.

c. Find and sketch a disc in \mathbb{R}^2 which you are sure contains a root.

2.26 There are other plausible ways to measure matrices other than the length and the norm; for example, we could declare the size $|A|$ of a matrix A to be the largest absolute value of an entry. In this case, $|A + B| \leq |A| + |B|$, but the statement $|A\vec{x}| \leq |A||\vec{x}|$ (where $|\vec{x}|$ is the ordinary length of a vector) is false. Find an ϵ so that it is false for

$$A = \begin{bmatrix} 1 & 1 & 1 + \epsilon \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \text{and} \quad \vec{x} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

2.27 Show that $\|A\| = \|A^\top\|$.

2.28 In example 2.10.9 we found that $M = 2\sqrt{2}$ is a global Lipschitz ratio for the function $\mathbf{f}\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} \sin(x+y) \\ x^2 - y^2 \end{pmatrix}$. What Lipschitz ratio do you get using the method of second partial derivatives? Using that Lipschitz ratio, what minimum domain do you get for the inverse function at $\begin{pmatrix} 0 \\ \pi \end{pmatrix}$?

Exercise 2.23: Note that

$$[2I]^3 = [8I], \quad \text{i.e.,}$$

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}^3 = \begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 8 \end{bmatrix}.$$

Exercise 2.24: The computation really does require you to row reduce a 4×4 matrix.

$$F\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} x^2 - y - 2 \\ y^2 - x - 6 \end{pmatrix}$$

Mapping F for exercise 2.25

Exercise 2.27: The norm $\|A\|$ of a matrix A is defined in section 2.9 (definition 2.9.6).

2.29 a. True or false? The equation $\sin(xyz) = z$ expresses x implicitly as a differentiable function of y and z near the point $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \pi/2 \\ 1 \\ 1 \end{pmatrix}$.

Exercise 2.30: You may use the fact that if

$$S : \text{Mat}(2, 2) \rightarrow \text{Mat}(2, 2)$$

is the squaring map

$$S(A) = A^2,$$

then

$$[\mathbf{D}S(A)]B = AB + BA.$$

b. True or false? The equation $\sin(xyz) = z$ expresses z implicitly as a differentiable function of x and y near the same point.

2.30 True or false? There exists a neighborhood $U \subset \text{Mat}(2, 2)$ of $\begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix}$ and a C^1 mapping $F : U \rightarrow \text{Mat}(2, 2)$ with

$$F\left(\begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}\right) = \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix}, \quad \text{and} \quad (F(A))^2 = A.$$

2.31 True or false? (Explain your answer.) There exists $r > 0$ and a differentiable map $g : B_r\left(\begin{bmatrix} -3 & 0 \\ 0 & -3 \end{bmatrix}\right) \rightarrow \text{Mat}(2, 2)$ such that

$$g\left(\begin{bmatrix} -3 & 0 \\ 0 & -3 \end{bmatrix}\right) = \begin{bmatrix} 1 & 2 \\ -2 & -1 \end{bmatrix}$$

$$\text{and } (g(A))^2 = A \text{ for all } A \in B_r\left(\begin{bmatrix} -3 & 0 \\ 0 & -3 \end{bmatrix}\right).$$

Dierk Schleicher contributed exercise 2.32. Geometrically, the condition given is the condition that there exists a unit cube with a vertex at the origin such that the three sides emanating from the origin are $\vec{\mathbf{a}}$, $\vec{\mathbf{b}}$, and $\vec{\mathbf{c}}$.

2.32 Given three vectors $\begin{bmatrix} a_1 \\ a_2 \end{bmatrix}, \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}, \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$ in \mathbb{R}^2 , show that there exist vectors $\vec{\mathbf{a}} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}, \vec{\mathbf{b}} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}, \vec{\mathbf{c}} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$ in \mathbb{R}^3 such that

$$|\vec{\mathbf{a}}|^2 = |\vec{\mathbf{b}}|^2 = |\vec{\mathbf{c}}|^2 = 1 \quad \text{and} \quad \vec{\mathbf{a}} \cdot \vec{\mathbf{b}} = \vec{\mathbf{a}} \cdot \vec{\mathbf{c}} = \vec{\mathbf{b}} \cdot \vec{\mathbf{c}} = 0$$

if and only if $\vec{\mathbf{v}}_1 = \begin{bmatrix} a_1 \\ b_1 \\ c_1 \end{bmatrix}$ and $\vec{\mathbf{v}}_2 = \begin{bmatrix} a_2 \\ b_2 \\ c_2 \end{bmatrix}$ are of unit length and orthogonal.

2.33 Imagine that, when constructing a Newton sequence

$$\mathbf{x}_{n+1} = \mathbf{x}_n - [\mathbf{D}\mathbf{f}(\mathbf{x}_n)]^{-1}\mathbf{f}(\mathbf{x}_n),$$

you happen upon a noninvertible matrix $[\mathbf{D}\mathbf{f}(\mathbf{x}_n)]$. What should you do? Suggest ways to deal with the situation.

2.34 Let V be a vector space, and consider the set of nonempty subsets of V , denoted $\mathcal{P}^*(V)$. Define $+ : \mathcal{P}^*(V) \times \mathcal{P}^*(V) \rightarrow \mathcal{P}^*(V)$ by

$$A + B := \{a + b \mid a \in A, b \in B\}$$

and scalar multiplication $\mathbb{R} \times \mathcal{P}^*(V) \rightarrow \mathcal{P}^*(V)$ by

$$\alpha A := \{\alpha a \mid a \in A\}.$$

a. Show that $+$ is associative: $(A + B) + C = A + (B + C)$ and that $\{0\}$ is a neutral element for $+$.